

Space-based diagnosis of surface ozone sensitivity to anthropogenic emissions

Randall V. Martin,^{1,2} Arlene M. Fiore,³ and Aaron Van Donkelaar¹

Received 5 January 2004; revised 17 February 2004; accepted 24 February 2004; published 24 March 2004.

[1] We present a novel capability in satellite remote sensing with implications for air pollution control strategy. We show that the ratio of formaldehyde columns to tropospheric nitrogen dioxide columns is an indicator of the relative sensitivity of surface ozone to emissions of nitrogen oxides ($\text{NO}_x \equiv \text{NO} + \text{NO}_2$) and volatile organic compounds (VOCs). The diagnosis from these space-based observations is highly consistent with current understanding of surface ozone chemistry based on in situ observations. The satellite-derived ratios indicate that surface ozone is more sensitive to emissions of NO_x than of VOCs throughout most continental regions of the Northern Hemisphere during summer. Exceptions include Los Angeles and industrial areas of Germany. A seasonal transition occurs in the fall when surface ozone becomes less sensitive to NO_x and more sensitive to VOCs.

INDEX TERMS: 0322 Atmospheric Composition and Structure: Constituent sources and sinks; 0345 Atmospheric Composition and Structure: Pollution—urban and regional (0305); 0365 Atmospheric Composition and Structure: Troposphere—composition and chemistry; 0368 Atmospheric Composition and Structure: Troposphere—constituent transport and chemistry; 0394 Atmospheric Composition and Structure: Instruments and techniques. **Citation:** Martin, R. V., A. M. Fiore, and A. Van Donkelaar (2004), Space-based diagnosis of surface ozone sensitivity to anthropogenic emissions, *Geophys. Res. Lett.*, 31, L06120, doi:10.1029/2004GL019416.

1. Introduction

[2] Surface ozone is deleterious to human health, crops, and ecosystems [*National Research Council*, 1991]. Uncertainty in the relation between surface ozone and its two main precursors, nitrogen oxides ($\text{NO}_x \equiv \text{NO} + \text{NO}_2$) and volatile organic compounds (VOCs) remains a primary obstacle to improving surface air quality [*Sillman*, 1999]. Ozone is produced in both NO_x -sensitive and NO_x -saturated (also known as VOC-sensitive) regimes. The design of control strategies for surface ozone has been impeded by limited observations of ozone- NO_x -VOC sensitivity [*Sillman*, 1999]. We develop a method to use space-based observations from the Global Ozone Monitoring Experiment (GOME) [*European Space Agency*, 1995; *Burrows et al.*, 1999] to characterize ozone- NO_x -VOC sensitivity in

surface air and apply our method to determine the spatial and temporal variation in ozone- NO_x -VOC sensitivity throughout the Northern Hemisphere.

[3] Surface ozone is produced by a chain reaction involving the photochemical oxidation of VOCs in the presence of NO_x . The chain reaction is propagated by cycling of HO_x ($\equiv \text{OH} + \text{peroxy}$) radicals. The pathway for HO_x loss is determined by the relative abundance of HO_x and NO_x . Ozone production tends to be either NO_x -sensitive if HO_x -loss occurs primarily by self reaction of peroxy radicals, or NO_x -saturated if the primary HO_x -loss pathway is via reaction of nitrogen dioxide (NO_2) and OH [*Sillman et al.*, 1990]. *Sillman* [1995] first presented the concept of using ozone- NO_x -VOC indicators to diagnose NO_x -sensitive versus NO_x -saturated conditions, and showed that the ratio of formaldehyde (HCHO) to total reactive nitrogen is such an indicator. A related ratio can be retrieved from the GOME satellite instrument, specifically the ratio of the HCHO column to the tropospheric NO_2 column.

2. Evaluation of the HCHO/ NO_2 Column Ratio as an Ozone- NO_x -VOC Indicator

[4] We use the GEOS-CHEM global 3-D model of ozone- NO_x -VOC chemistry [*Bey et al.*, 2001], version 4.16 [*Martin et al.*, 2002a], to evaluate the quality of the HCHO/ NO_2 column ratio as an indicator of NO_x -sensitive versus NO_x -saturated conditions. We analyze results from three simulations previously described by *Fiore et al.* [2002]: a standard simulation, a simulation in which anthropogenic VOC emissions are reduced by 50%, and a simulation in which anthropogenic NO_x emissions are reduced by 50%. The simulations use assimilated observations of meteorological fields from NASA GEOS-1 with 20 vertical levels and a $4^\circ \times 5^\circ$ horizontal resolution. All simulations are spun up for 10 months to remove the effects of initial conditions. We present results for May–November.

[5] Figure 1 shows the calculated response in monthly-mean afternoon surface ozone over polluted regions to a prescribed change in emissions as a function of the tropospheric HCHO/ NO_2 column ratio. The transition between NO_x -sensitive and NO_x -saturated regimes occurs for a HCHO/ NO_2 ratio of about 1. HCHO/ NO_2 ratios above 1 tend to be NO_x -sensitive as indicated by the decrease in surface ozone for the simulation with reduced NO_x emissions. HCHO/ NO_2 ratios below 1 reflect NO_x -saturated conditions; reductions in NO_x emissions increase surface ozone while reductions in VOC emissions decrease surface ozone.

[6] Why does the HCHO/ NO_2 column ratio function effectively as an indicator of surface ozone- NO_x -VOC sensitivity? First, the bulk of the tropospheric NO_2 and

¹Department of Physics and Atmospheric Science, Dalhousie University, Halifax, Nova Scotia, Canada.

²Harvard-Smithsonian Center for Astrophysics, Cambridge, Massachusetts, USA.

³Atmospheric and Oceanic Sciences Program, Princeton University, Princeton, New Jersey, USA.




Figure 1. Model calculation of the normalized local sensitivity of afternoon surface ozone, $d[\text{O}_3]/dE$, to a 50% reduction in anthropogenic emissions, dE , in units of either molecules nitrogen $\text{m}^{-2} \text{s}^{-1}$ or molecules carbon $\text{m}^{-2} \text{s}^{-1}$. The local sensitivity and local model tropospheric HCHO/NO₂ column ratio are monthly mean values from May to November. Afternoon surface ozone is the average between 1 pm and 5 pm local time. The calculation is restricted to polluted regions (model tropospheric NO₂ column $>2.5 \times 10^{15}$ molecules cm^{-2}).

HCHO columns over polluted regions is within the mixed layer [Ladstätter-Weissenmayer *et al.*, 2003]. Second, the division between NO_x-sensitive and NO_x-saturated regimes generally follows a constant reactivity-weighted VOC/NO_x ratio [Chameides *et al.*, 1992], a ratio closely associated to the HCHO/NO₂ ratio. Third, the HCHO/NO₂ column ratio largely reflects the ratio of HO_x to NO_x sources near the surface. HCHO is positively correlated with the source of HO_x in polluted environments [Martinez *et al.*, 2003], and tropospheric NO₂ is closely related to NO_x emissions [Leue *et al.*, 2001; Martin *et al.*, 2003]. The ratio of the HO_x source to the NO_x source determines NO_x-VOC sensitivity, and the transition from NO_x-sensitive to NO_x-saturated regimes occurs at a specific ratio over a range of conditions [Kleinman, 1991; Jacob *et al.*, 1995].

3. Space-Based Observations of Tropospheric NO₂ and HCHO

[7] The nadir-viewing GOME satellite instrument has provided the capability for continuous global monitoring of tropospheric NO₂ and HCHO columns through observation of solar backscatter since 1995. Global coverage is achieved every 3 days after 43 orbits with a typical surface spatial resolution of 40 km by 320 km, sufficient to resolve the regional scale of ozone pollution episodes [Logan, 1989]. Observations at northern mid-latitudes occur between 10:30 AM–11:30 AM local solar time. Our numerical model calculations show that the HCHO/NO₂ ratio exhibits little diurnal variation during daytime; the ratio at 11 AM is typically within 5% of the ratio in the afternoon, so the thresholds identified in Figure 1 are applicable to GOME observations.

[8] The GOME retrievals are described in Martin *et al.* [2002b, 2003] for tropospheric NO₂ and in Chance *et al.* [2000] and Abbot *et al.* [2003] for HCHO. These retrievals include the air mass factor (AMF) formulation of Palmer *et al.* [2001] and radiative transfer calculations with the Linearized Discrete Ordinate Radiative Transfer (LIDORT) model developed by Spurr [2002]. Observation scenes are

excluded when local cloud cover retrieved from GOME [Kurosu *et al.*, 1999] exceeds 40%.

[9] Errors in the GOME HCHO/NO₂ ratio are reduced by compensating effects. The largest source of error in individual trace gas retrievals is from the AMF calculation due to surface reflectivity, clouds, aerosols, and trace gas profile [Martin *et al.*, 2002b; Boersma *et al.*, 2004]. Error in the AMF calculation similarly affects NO₂ and HCHO retrievals and largely cancels in the ratio. Uncertainty in the resulting monthly mean ratio is typically less than 25%.

4. Ozone-NO_x-VOC Sensitivity Diagnosed From Space-Based Observations

[10] Figure 2 shows the GOME HCHO/NO₂ column ratio over North America, Europe, and East Asia during May to November 1997. Ratios greater than one, indicating NO_x-sensitive conditions over most suburban and rural areas during summer, are consistent with analysis of in situ observations at rural sites of North America by Trainer *et al.* [1993]. The decrease in the HCHO/NO₂ ratio to less than one over the eastern United States during fall shows the seasonal transition from NO_x-sensitive to NO_x-saturated conditions due to a decline in isoprene and HO_x that is expected from theory [Kleinman, 1991], in situ [Jacob *et al.*, 1995], and models [Liang *et al.*, 1998]. The seasonal decline in isoprene and HO_x is also associated with a seasonal decline in the HCHO column to values below the GOME detection limit.

[11] GOME shows that major urban and industrial centers such as Los Angeles tend to be NO_x-saturated, but transition to a NO_x-sensitive regime downwind, consistent with model calculations by Milford *et al.* [1989]. The initial NO_x supply greatly exceeds the HO_x supply in major urban areas, but a switch from NO_x-saturated to NO_x-sensitive conditions occurs from the combined effects of dilution and HO_x production downwind, as the HO_x supply surpasses the initial NO_x source [Kleinman, 1991, 1994].

[12] Data over Europe are sparse due to HCHO columns below the GOME detection limit. London is near NO_x-saturation, consistent with calculations of Derwent *et al.*




Figure 2. Monthly mean tropospheric HCHO/NO₂ column ratio retrieved from the GOME satellite instrument for North America, Europe, and East Asia. Ratios greater than 1 tend to be NO_x-sensitive. Ratios below 1 tend to be NO_x-saturated. White areas indicate remote regions (observed tropospheric NO₂ columns less than 2.5×10^{15} molecules cm⁻²) and regions below the HCHO detection limit of 4×10^{15} molecules cm⁻².

[2003]. In situ measurements [Staffelbach *et al.*, 1997] support GOME observations that southern Switzerland is NO_x-sensitive during summer. GOME reveals that industrial Germany is NO_x-saturated throughout the year, and that a transition from NO_x-sensitive to NO_x-saturated conditions occurs over the Po Valley from September to October.

[13] GOME HCHO/NO₂ ratios greater than one during summer over eastern China, Korea, and Japan indicate NO_x-

sensitive conditions. However, there is a clear transition to NO_x-saturated conditions in the northern regions during fall. The resulting gradient during October and November of NO_x-sensitive conditions in rural southern China, but NO_x-saturated conditions elsewhere, is consistent with photochemical model calculations of Luo *et al.* [2000]. This transition is of particular interest since episodes of enhanced ozone concentrations are frequent in the fall, and may be reducing crop yields [Chameides *et al.*, 1999].

[14] Results presented here suggest that nascent capabilities in satellite remote sensing are of sufficient quality to contribute to air pollution management by providing guidance on the spatial and temporal sensitivity of surface ozone to reductions of NO_x and VOC emissions. Such observations also provide a critical test of air quality models used to develop emission controls. These observational capabilities will be enhanced by the next generation of instruments, such as the SCIAMACHY and OMI satellite instruments, which feature higher spatial resolution that should resolve transitions between urban and suburban environments. Future work should systematically examine the relationship of these space-based column indicators with high-resolution simulations and in situ indicators.

[15] **Acknowledgments.** We thank Ian Folkins, Daniel Jacob, and two anonymous reviewers for helpful comments that contributed to substantial improvement to the manuscript. We are grateful to Hongyu Liu for providing meteorological fields. This work was supported by Dalhousie University, by NASA's Radiation Science Program, and by the Postdoctoral Research Program in Atmospheric and Oceanic Sciences at Princeton University and GFDL/NOAA. The GEOS-CHEM model is managed by the Atmospheric Chemistry Modeling Group at Harvard University with support from the NASA Atmospheric Chemistry Modeling and Analysis Program.

References

- Abbot, D. S., P. I. Palmer, R. V. Martin, K. Chance, D. J. Jacob, and A. Guenther (2003), Seasonal and interannual variability of North American isoprene emissions as determined by formaldehyde column measurements from space, *Geophys. Res. Lett.*, *30*(17), 1886, doi:10.1029/2003GL017336.
- Bey, I., et al. (2001), Global modeling of tropospheric chemistry with assimilated meteorology: Model description and evaluation, *J. Geophys. Res.*, *106*, 23,073–23,096.
- Boersma, K. F., H. J. Eskes, and E. J. Brinksma (2004), Error analysis for tropospheric NO₂ retrieval from space, *J. Geophys. Res.*, *109*, D04311, doi:10.1029/2003JD003962.
- Burrows, J. P., et al. (1999), The Global Ozone Monitoring Experiment (GOME): Mission concept and first scientific results, *J. Atmos. Sci.*, *56*, 151–175.
- Chameides, W. L., et al. (1992), Ozone precursor relationships in the ambient atmosphere, *J. Geophys. Res.*, *97*, 6037–6056.
- Chameides, W. L., et al. (1999), Is ozone pollution affecting crop yields in China?, *J. Geophys. Res.*, *26*, 867–870.
- Chance, K., P. I. Palmer, R. J. D. Spurr, R. V. Martin, T. P. Kurosu, and D. J. Jacob (2000), Satellite observations of formaldehyde over North America from GOME, *Geophys. Res. Lett.*, *27*, 3461–3464.
- Derwent, R. G., et al. (2003), Photochemical ozone formation in north west Europe and its control, *Atmos. Environ.*, *37*, 1983–1991.
- European Space Agency (1995), *The GOME Users Manual*, edited by F. Bednarz, Publ. SP-1182, Noordwijk, Netherlands.
- Fiore, A. M., D. J. Jacob, B. D. Field, D. G. Streets, S. D. Fernandes, and C. Jang (2002), Linking ozone pollution and climate change: The case for controlling methane, *Geophys. Res. Lett.*, *29*(19), 1919, doi:10.1029/2002GL015601.
- Jacob, D. J., L. W. Horowitz, J. W. Munger, B. G. Heikes, R. R. Dickerson, R. S. Artz, and W. C. Keene (1995), Seasonal transition from NO_x- to hydrocarbon-limited conditions for ozone production over the eastern United States in September, *J. Geophys. Res.*, *100*, 9315–9324.
- Kleinman, L. I. (1991), Seasonal dependence of boundary layer peroxide concentration: The low and high NO_x regimes, *J. Geophys. Res.*, *96*, 20,721–20,733.
- Kleinman, L. I. (1994), Low and high NO_x tropospheric photochemistry, *J. Geophys. Res.*, *99*, 16,831–16,838.
- Kurosu, T. P., K. Chance, and R. J. D. Spurr (1999), CRAG—Cloud Retrieval Algorithm for ESA's Global Ozone Monitoring Experiment, paper presented at the European Symposium on Atmospheric Measurements From Space, Eur. Space Agency Space Sci. Div., Noordwijk, Netherlands, 18–22 Jan.
- Ladstätter-Weissenmayer, A., et al. (2003), Transport and build-up of tropospheric trace gases during the MINOS campaign: Comparison of GOME, in situ aircraft measurements and MATCH-MPIC-data, *Atmos. Chem. Phys.*, *3*, 1887–1902.
- Leue, C., M. Wenig, T. Wagner, O. Klimm, U. Platt, and B. Jähne (2001), Quantitative analysis of NO_x emissions from Global Ozone Monitoring Experiment satellite image sequences, *J. Geophys. Res.*, *106*, 5493–5505.
- Liang, J., L. W. Horowitz, D. J. Jacob, Y. Wang, A. M. Fiore, J. A. Logan, G. M. Gardner, and J. W. Munger (1998), Seasonal budgets of reactive nitrogen species and ozone over the United States, and export fluxes to the global atmosphere, *J. Geophys. Res.*, *103*, 13,435–13,450.
- Logan, J. A. (1989), Ozone in rural areas of the United States, *J. Geophys. Res.*, *94*, 8511–8532.
- Luo, C., J. C. St. John, Z. Xiuji, K. S. Lam, T. Wang, and W. L. Chameides (2000), A nonurban ozone air pollution episode over eastern China: Observations and model simulations, *J. Geophys. Res.*, *105*, 1889–1908.
- Martin, R. V., et al. (2002a), Interpretation of TOMS observations of tropical tropospheric ozone with a global model and in situ observations, *J. Geophys. Res.*, *107*(D18), 4351, doi:10.1029/2001JD001480.
- Martin, R. V., et al. (2002b), An improved retrieval of tropospheric nitrogen dioxide from GOME, *J. Geophys. Res.*, *107*(D20), 4437, doi:10.1029/2001JD001027.
- Martin, R. V., D. J. Jacob, K. Chance, T. P. Kurosu, P. I. Palmer, and M. J. Evans (2003), Global inventory of nitrogen oxide emissions constrained by space-based observations of NO₂ columns, *J. Geophys. Res.*, *108*(D17), 4537, doi:10.1029/2003JD003453.
- Martinez, M., et al. (2003), OH and HO₂ concentrations, sources, and loss rates during the Southern Oxidants Study in Nashville, Tennessee, summer 1999, *J. Geophys. Res.*, *108*(D19), 4617, doi:10.1029/2003JD003551.
- Milford, J. B., A. G. Russell, and G. J. McRae (1989), A new approach to photochemical pollution control: Implications of spatial patterns in pollutant responses to reductions in nitrogen oxides and reactive organic gas emissions, *Environ. Sci. Technol.*, *23*, 1290–1301.
- National Research Council (1991), *Rethinking the Ozone Problem in Urban and Regional Air Pollution*, Nat. Acad. Press, Washington, D. C.
- Palmer, P. I., et al. (2001), Air mass factor formulation for spectroscopic measurements from satellites: Application to formaldehyde retrievals from the Global Ozone Monitoring Experiment, *J. Geophys. Res.*, *106*, 14,539–14,550.
- Sillman, S. (1995), The use of NO₂, H₂O₂, and HNO₃ as empirical indicators for ozone-NO_x-hydrocarbon sensitivity in urban locations, *J. Geophys. Res.*, *100*, 14,175–14,188.
- Sillman, S. (1999), The relation between ozone, NO_x, and hydrocarbons in urban and polluted rural environments, *Atmos. Environ.*, *33*, 1821–1845.
- Sillman, S., J. A. Logan, and S. C. Wofsy (1990), The sensitivity of ozone to nitrogen oxides and hydrocarbons in regional ozone episodes, *J. Geophys. Res.*, *95*, 1837–1852.
- Spurr, R. J. D. (2002), Simultaneous derivation of intensities and weighting functions in a general pseudo-spherical discrete ordinate radiative transfer treatment, *J. Quant. Spectrosc. Radiat. Transfer*, *75*, 129–175.
- Staffelbach, T., et al. (1997), Photochemical oxidant formation over southern Switzerland: 1. Results from summer 1994, *J. Geophys. Res.*, *102*, 23,345–23,362.
- Trainer, M., et al. (1993), Correlation of ozone with NO_y in photochemically aged air, *J. Geophys. Res.*, *98*, 2917–2925.

A. Van Donkelaar and R. V. Martin, Department of Physics and Atmospheric Science, Dalhousie University, Halifax, Nova Scotia, Canada B3H 3J5. (rvmartin@fizz.phys.dal.ca)

A. M. Fiore, Atmospheric and Oceanic Sciences Program, Princeton University, Princeton, NJ 08544, USA.